Supplementary Material

Article Title: Childhood Trauma Is Associated With Severe Clinical Characteristics of Bipolar Disorders

Author(s): Bruno Etain, MD, PhD; Monica Aas, PhD; Ole A. Andreassen, MD, PhD; Steinar Lorentzen, MD, PhD; Ingrid Dieset, MD; Sebastien Gard, MD; Jean-Pierre Kahn, MD; Frank Bellivier, MD, PhD; Marion Leboyer, MD, PhD; Ingrid Melle, MD, PhD; and Chantal Henry, MD

List of Supplementary Material for the article

Disclaimer

This Supplementary Material has been provided by the author(s) as an enhancement to the published article. It has been approved by peer review; however, it has undergone neither editing nor formatting by in-house editorial staff. The material is presented in the manner supplied by the author.

Supplementary eTable 1. Demographics comparison between sites

	French site	Norwegian site	Statistics
Age (mean±SD)	43.50±13.24	33.25±11.67	t=8.78, df=585, p<0.001
Gender, M/F	168/250	66/103	X^2=0.07, df=1, p=0.80
Suicide attempts (yes/no)	156/256	55/112	X^2=1.25, df=1, p=0.26
Depressive episodes (mean±SD)	4.57±5.29	5.36±6.02	t=-1.44, df=275, p=0.15
Manic/hypomanic episodes (mean±SD)	4.72±6.05	4.35±5.98	t=0.66, df=554, p=0.51
Psychotic episodes (Yes/No)	163/181	84/72	X^2=1.79, df=1, p=0.18
Rapid Cycling (Yes/No)	89/297	25/136	X^2=3.90, df=1, p=0.05
Lifetime cannabis abuse (Yes/No)	50/362	25/144	X^2=0.75, df=1, p=0.39
Lifetime alcohol dependence (Yes/No)	30/382	23/146	X^2=5.79, df=1, p=0.016
Age at onset (mean±SD)	25.17±10.32	21.87±9.24	t=3.59, df=577, p<0.001
Duration of illness (mean±SD)	18.36±12.41	11.39±10.03	t=7.06, df=377, p<0.001
CTQ total score (mean±SD)	41.75±12.89	44.03±16.43	t=-1.57, df=237, p=0.12
Diagnosis			X^2=0.61, df=2, p=0.74
Bipolar 1, n	306	119	
Bipolar 2, n	88	38	
Bipolar NOS, n	24	12	

CTQ=Childhood Trauma Questionnaire, F=female, M=male, NOS=not otherwise specified.