

Emotion Dysregulation and Negative Affect: Association With Psychiatric Symptoms

Bekh Bradley, PhD; Jared A. DeFife, PhD; Clifford Guarnaccia, PhD;
Justine Phifer, BA; Negar Fani, MA; Kerry J. Ressler, MD, PhD; and Drew Westen, PhD

Objective: A growing body of research focuses on the development and correlates of emotion dysregulation, or deficits in the ability to regulate intense and shifting emotional states. Current models of psychopathology have incorporated the construct of emotion dysregulation, suggesting its unique and interactive contributions, along with childhood disruptive experiences and negative affect, in producing symptomatic distress. Some researchers have suggested that emotion dysregulation is simply a variant of high negative affect. The aim of this study was to assess the construct and incremental validity of self-reported emotion dysregulation over and above childhood trauma and negative affect in predicting a range of psychopathology.

Method: Five hundred thirty individuals aged 18 to 77 years (62% female) were recruited from the waiting areas of the general medical and obstetric/gynecologic clinics in an urban public hospital in Atlanta, Georgia. Participants completed a battery of self-report measures obtained by interview, including the Childhood Trauma Questionnaire, the Positive and Negative Affect Schedule, and the Emotion Dysregulation Scale. Regression analyses examined the unique and incremental associations of these self-report measurements of childhood traumatic experiences, negative affect, and emotion dysregulation with concurrent structured interview-based measurements of psychiatric distress and history of self-destructive behaviors. These measures included the Clinician-Administered PTSD Scale, the Alcohol Use Disorders Identification Test, the Short Drug Abuse Screening Test, the Beck Depression Inventory, and the Global Adaptive Functioning Scale from the Longitudinal Interval Follow-Up Evaluation. The presented data were collected between 2005 and 2009.

Results: Regression models including age, gender, childhood trauma, negative affect, and emotion dysregulation were significantly ($P \leq .001$) associated with each of the study's criterion variables, accounting for large portions of the variance in posttraumatic stress symptoms ($R^2 = 0.21$), alcohol and drug abuse ($R^2 = 0.28$ and 0.21 , respectively), depression ($R^2 = 0.55$), adaptive functioning ($R^2 = 0.14$), and suicide history (omnibus $\chi^2 = 74.80$, $P < .001$). Emotion dysregulation added statistically significant ($P < .01$) incremental validity to each regression model ($\beta = 0.25, 0.34, 0.35, 0.34$, and -0.18 , and Wald = 24.43, respectively).

Conclusions: Results support the conceptualization of emotion dysregulation as a distinct and clinically meaningful construct associated with psychiatric distress that is not reducible to negative affect. Emotion dysregulation is a key component in a range of psychiatric symptoms and disorders and a core target for psychopharmacologic and psychosocial treatment interventions.

J Clin Psychiatry 2011;72(5):685–691

© Copyright 2011 Physicians Postgraduate Press, Inc.

Submitted: July 12, 2010; accepted August 20, 2010

(doi:10.4088/JCP.10m06409blu).

Corresponding author: Jared A. DeFife, PhD, Laboratory of Personality and Psychopathology, Department of Psychology, Emory University, 36 Eagle Row, Ste 415, Atlanta, GA 30322 (jdefife@emory.edu).

A well-established body of literature has identified a trait-like tendency toward negative affect or “neuroticism” as a higher order construct central to multiple Axis I and Axis II disorders.^{1,2} In addition, deficits in the ability to regulate intense, negative, and shifting emotional states may be implicated in various forms of adult psychopathology. As such, accumulating of research has focused on the defining properties, developmental trajectory, neurobiological underpinnings, and behavioral manifestations of emotion dysregulation.³

Components of emotion dysregulation include a tendency for emotions to spiral out of control, change rapidly, get expressed in intense and unmodified forms, and/or overwhelm both coping capacity and reasoning.^{4–8} Developmental research suggests that these self-regulatory deficits emerge from an interaction of intrinsic temperamental and biological factors as well as extrinsic intrusions from chaotic and stressful early life experiences, particularly childhood abuse and problematic attachments with caregivers.⁹ This model suggests that children prone to negative affect and/or higher levels of emotional reactivity will be at higher risk for emotion dysregulation when exposed to chaotic, disruptive environments and other adverse childhood events. The strong association between adverse childhood experiences and a wide range of adult mental problems (eg, depression, suicide, substance use) may be, in part, accounted for by vulnerability to poor emotion regulation among children exposed to multiple adverse childhood events.^{10–14}

A small but growing number of studies have begun to focus on both negative affect (ie, emotions people tend to feel) and emotion regulation (ie, the ability to adaptively manage emotions, particularly negative emotions). The prevailing focus of research on emotion dysregulation in clinical populations has centered on borderline personality disorder, identifying both negative affect and emotion dysregulation as central to this form of psychopathology.^{15–20} Cross-sectional and longitudinal studies^{21–25} utilizing various self-report and expert-report measures have found that emotion dysregulation is a highly descriptive, distinctive, temporally stable, and heritable component of borderline personality disorder.^{21–25} Even after analyses were controlled for levels of depression, emotional intensity and dysregulation related significantly to a number of borderline personality disorder traits.²⁶

FOR CLINICAL USE

- ◆ Emotion dysregulation is inherent in a number of forms of psychopathology and may present in extremes of either overly restricted emotional expression and avoidance or heightened and excessive emotionality and excitement-seeking.
- ◆ Treatment goals for individuals with emotion dysregulation may include reduction of affective arousal, increase in affect tolerance, and development of psychosocial coping skills to be used in times of increased emotional distress.
- ◆ Techniques focusing on social-perspective-taking and cognitive reappraisal of negative affect states, causes, and consequences may be beneficial for individuals with emotion dysregulation.

Similarly, comparisons of patients with dysthymia to those with borderline personality disorder find that dysthymic patients are characterized by negative affect alone, whereas borderline personality disorder patients are characterized by both negative affect and emotion dysregulation.^{15,27} Both emotion dysregulation and negative affect are primary targets of a number of treatment approaches for borderline personality disorder.^{28–31}

Across Axis II syndromes, neuroticism (defined as a predisposition to negative affective states) demonstrates a distinctive associative pattern with cluster C personality disorders as compared to the affective instability associated with cluster B disorders.³² In a study of patients receiving treatment for substance dependence, McDermott and colleagues³³ found that emotion regulation difficulties distinguished posttraumatic stress disorder (PTSD) patients from non-PTSD patients above and beyond the level of anxiety symptom severity.

The goal of this study was to examine the relationship of both negative affect and emotion dysregulation with symptoms of multiple Axis I symptoms including depression, PTSD symptoms, alcohol and substance use-related symptoms, and history of suicide attempts, as well as global adaptive functioning. Specifically, we examined the contributions of emotion dysregulation, childhood trauma, and negative affect to a range of clinical psychopathology.

We are focusing on these variables as they present in a population at high risk of vulnerability to psychiatric disorders including depression, PTSD, and substance abuse.^{34–36} Understanding the factors underlying risk and resilience to mental disorders in this population is a question that is of high public health importance. Using the self-report Emotion Dysregulation Scale, the Childhood Trauma Questionnaire, and the Positive and Negative Affect Scale, we examined selected a priori criterion variables of clinical interest including measurements of posttraumatic stress, substance abuse history, depressive symptoms, suicide attempt history, and global adaptive functioning.

METHOD

Procedure

Subjects in this study were ascertained as part of the Grady Trauma Project, a 5-year National Institutes of Health-funded study of risk and resilience factors related to

PTSD.^{34,37,38} Participants were recruited from the general medical and obstetric/gynecologic clinics at a publicly funded, not-for-profit health care system that serves a low-income population in Atlanta, Georgia. Interviewers approached participants waiting for appointments. Participants were read each question by a trained interviewer who recorded their responses onto a tablet computer. Participants completed a battery of self-report measures that took 45–75 minutes to complete (dependent in large part on the extent of the participant's trauma history and symptoms). All measures were obtained by verbal interview. Each person was paid \$15.00 for participation in this phase of the study. Eligibility requirements for all phases of the study included the ability to give informed consent, and written and verbal informed consent was obtained for all participants. All procedures in this study were approved by the institutional review boards of Emory University School of Medicine and Grady Memorial Hospital, Atlanta, Georgia. Presented data were collected between 2005 and 2009.

As described in full detail previously,³⁴ study participants who completed this initial interview were invited to participate in a secondary phase of the study that included a more comprehensive, structured interview-based assessment of psychological functioning.^{34,38}

Participants

The sample included 530 study participants (although, as some participants did not complete all measures, the number of participants for individual analyses varies). The participants were predominantly female (62%), with ages ranging from 18 to 77 years (mean = 42.3 years, SD = 12.6 years). Eighty-eight percent of participants were African American, 5% were white, 1% were Latino, and 3% were mixed or other. Twenty-four percent had less than a 12th-grade education, 42% completed a terminal high school diploma or General Educational Development test, 22% had some college or technical school education, and 8% graduated from a college or technical school.

Measures

Childhood Trauma Questionnaire. The Childhood Trauma Questionnaire^{39,40} is a 28-item self-report measure of childhood trauma and neglect assessing 5 types of maltreatment: sexual, physical, and emotional abuse and emotional and physical neglect.

Positive and Negative Affect Schedule. The Positive and Negative Affect Schedule⁴¹ is a well-validated brief measure of general mood state. Participants were asked to rate on a 5-point Likert scale their general experiences with 20 emotion adjectives, 10 describing positive emotional states (eg, excited, proud, and inspired) and 10 describing negative emotional states (eg, distressed, jittery, and irritable). Analyses for this study focused on the negative affect portion of the scale only.

Emotion Dysregulation Scale. The Emotion Dysregulation Scale (D.W., B.B., unpublished scale, 2008; available at <http://www.psychsystems.net>) is a 24-item self-report scale adapted from the clinician-rated Affect Regulation and Experience Q-Sort Questionnaire.^{15,27,42,43} Items are scored on a 7-point Likert scale and assess domains of emotional experiencing (eg, "My emotions sometimes spiral out of control," "Emotions overwhelm me," "When I feel angry, I get really angry"); cognition (eg, "When I'm upset, I have trouble seeing or remembering anything good about myself," "When I'm feeling bad, I have trouble remembering anything positive; everything just feels bad"); and behavior (eg, "When my emotions are strong, I often make bad decisions," "When I'm upset, I sometimes become needy or clingy"). The internal consistency of the scale is high ($\alpha = 0.97$).

Clinician-Administered PTSD Scale. The Clinician-Administered PTSD Scale⁴⁴ is an interviewer-administered diagnostic instrument measuring PTSD, and it includes items that rate social and occupational functioning, global PTSD symptom severity, and the validity of the participant's responses. The Clinician-Administered PTSD Scale assesses PTSD diagnosis and yields a continuous measure of the severity of overall PTSD and its 3 symptom clusters (intrusion, avoidance, and arousal). The frequency and intensity scores for each of the 17 diagnostic criteria are summed to arrive at a total severity score. This measure has excellent psychometric properties.^{45,46}

Alcohol Use Disorders Identification Test. The Alcohol Use Disorders Identification Test⁴⁷ is an interview-based assessment measuring frequency of both alcohol use and related behavioral problems, yielding a total score from 0 to 40.

Short Drug Abuse Screening Test. The Short Drug Abuse Screening Test⁴⁸ is a 10-item self-report measurement of non-alcohol substance use and related problems. Multiple studies support the psychometric properties of the Short Drug Abuse Screening Test in the assessment of drug abuse and dependence in a variety of settings and populations.⁴⁹⁻⁵¹

Beck Depression Inventory. The Beck Depression Inventory^{52,53} is a widely used, 21-item self-report measurement of depressive symptoms. In addition to the Beck Depression Inventory, participants were also asked to self-report any history of suicide attempts.

Longitudinal Interval Follow-Up Evaluation–Psychosocial Schedule–Global Adaptive Functioning Scale. A portion of the Longitudinal Interval Follow-Up Evaluation⁵⁴ interview was used to obtain a broad-based measurement of participants' subjective sense of global

adaptive functioning during the prior month. Participants were asked to rate their level of functioning during the prior month on a scale from 1 (very good, no impairment) to 5 (very poor, severe impairment).

RESULTS

To determine the extent of association between our predictor variables (childhood trauma, negative affect, and emotion dysregulation), we calculated Pearson correlation coefficients. Each correlation had a large effect size and was statistically significant. Childhood trauma and negative affect scores correlated at $r_{428} = 0.25$ ($P < .001$), while emotion dysregulation scores were also significantly related to childhood trauma ($r_{429} = 0.26$, $P < .001$) and negative affect ($r_{425} = 0.57$, $P < .001$).

We then conducted a series of linear regressions (for dimensional criterion variables) and a logistic regression (for the categorical criterion of reported suicide history) to examine the unique and combined associations of childhood trauma, negative affect, and emotion dysregulation ratings with variation in our selected criterion measurements of posttraumatic stress, substance abuse problems, depression, suicidality, and adaptive functioning. While the selected predictor variables were significantly correlated, the relationships were far from large enough in effect to create concerns about multicollinearity or variance inflation within a regression model.⁵⁵ In each regression, age and gender (coded as male = 0, female = 1) were entered in the first step of the model to control for demographic variations. We then entered childhood trauma and negative affect ratings in the second step and emotion dysregulation in the final step to examine the incremental validity of emotion dysregulation in each predictive model. Table 1 presents results from the third-step overall model of each linear regression, and Table 2 presents results of the logistic regression for suicide history.

Posttraumatic Stress

As seen in Table 1, an overall model including age, gender, childhood trauma, negative affect, and emotion dysregulation was significant ($F_{394} = 21.03$, $P \leq .001$), accounting for 21% of the variance in frequency and intensity of posttraumatic stress symptoms. Each of our selected predictor variables (childhood trauma, negative affect, and emotion dysregulation) was statistically significant, and adding emotional dysregulation in the final step accounted for a statistically significant incremental 4% ($P \leq .001$) of the overall variance (a unique R change of 0.16).

Substance Abuse

For alcohol use and related behavioral problems, the overall regression model was statistically significant ($F_{123} = 9.29$, $P \leq .001$), accounting for 28% of the variance in alcohol abuse. Age and gender were significantly related to reported alcohol abuse (Table 1), with older participants and male participants more likely to indicate problems stemming from alcohol abuse. Of our 3 target predictor variables, only

Table 1. Linear Regression Analyses of Predictor Variables With Measures of Psychological Functioning

	b	SE b	β	F	R ²
Posttraumatic stress disorder				21.03***	0.21
Age	0.07	0.12	0.03		
Gender	2.11	2.92	0.03		
Childhood trauma	0.36	0.08	0.21***		
Negative affect	0.49	0.19	0.14*		
Emotion dysregulation	0.19	0.04	0.25***		
Alcohol abuse				9.29***	0.28
Age	0.15	0.07	0.17*		
Gender	-7.88	1.68	-0.37***		
Childhood trauma	0.03	0.04	0.07		
Negative affect	-0.06	0.10	-0.05		
Emotion dysregulation	0.08	0.02	0.34***		
Drug abuse				5.06***	0.21
Age	0.05	0.02	0.22*		
Gender	-0.25	0.52	-0.04		
Childhood trauma	0.03	0.01	0.22*		
Negative affect	-0.06	0.03	-0.22		
Emotion dysregulation	0.02	0.01	0.35**		
Depression				96.72***	0.55
Age	0.04	0.04	0.04		
Gender	1.10	0.87	0.04		
Childhood trauma	0.09	0.02	0.14***		
Negative affect	0.62	0.06	0.44***		
Emotion dysregulation	0.10	0.01	0.34***		
Adaptive functioning				12.12***	0.14
Age	0.00	0.00	0.03		
Gender	0.20	0.10	0.10*		
Childhood trauma	-0.00	0.00	-0.07		
Negative affect	-0.02	0.01	-0.20***		
Emotion dysregulation	-0.00	0.00	-0.18**		

* $P < .05$, ** $P < .01$, *** $P \leq .001$.

emotion dysregulation was a significant predictor of alcohol abuse history, accounting for an 8% increment ($P \leq .001$) of the unique variance in the model.

The overall regression model for nonalcohol substance use and related problems was also statistically significant ($F_{103} = 5.06$, $P \leq .001$), accounting for 21% of the reported drug abuse variance. Age was significantly related to drug abuse problems, as were both childhood trauma and emotion dysregulation (Table 1). Again, adding emotion dysregulation to the final step of the regression model resulted in a statistically significant incremental prediction, accounting for an additional 7% of the unique variance ($P < .01$). Negative affect ratings demonstrated an inversely associated trend with drug abuse problems.

Depression and History of Suicide Attempt

The overall model predicting depressive symptoms was significant ($F_{394} = 96.72$, $P \leq .001$) and accounted for 55% of the variance in Beck Depression Inventory scores. Again, each of our hypothesized predictor variables was significantly related to depressive symptoms (Table 1), with emotion dysregulation accounting for an incremental 7% ($P < .01$) of the variance in the final prediction model.

Table 2 presents results from the final block of the logistic regression analysis for history of suicide attempt. The overall model was significant. In block 2 of the model, negative affect was significantly associated with suicide history (Wald statistic = 10.56, $P = .001$); however, this relationship was

Table 2. Logistic Regression Analysis of Predictor Variables With History of Suicide Attempt

	b	SE b	Exp(B)	Wald ^a
Suicide attempt				
Age	0.02	0.01	1.02	1.61
Gender	0.97	0.33	2.62	8.37**
Childhood trauma	0.03	0.01	1.03	15.24***
Negative affect	0.00	0.02	1.03	0.02
Emotion dysregulation	0.02	0.00	1.02	24.43***

^aOmnibus $\chi^2 = 74.80$, $P < .001$.

** $P < .01$, *** $P \leq .001$.

virtually eliminated when emotion dysregulation was added to the model in block 3. In the final model, female gender, childhood trauma, and emotion dysregulation scores were all significantly associated with suicide history.

Global Adaptive Functioning

As indicated in Table 1, the overall regression model for global adaptive functioning was statistically significant ($F_{386} = 12.12$, $P \leq .001$), although relatively smaller in effect, accounting for 14% of the variance. Gender was significantly related to higher self-reported ratings of global adaptive functioning in the prior month. Childhood trauma was not significantly related to recent adaptive functioning, while negative affect and emotion dysregulation were significantly related to lower adaptive functioning scores. Adding emotion dysregulation in the final step added a smaller but nevertheless significant increment to the model's predictive power, accounting for an additional 2% ($P < .01$) of the unique variance.

DISCUSSION

Our findings indicate that this self-report measure of emotion dysregulation was internally consistent and related, as expected, to a range of psychopathology criterion variables including posttraumatic stress, substance abuse problems, depression, suicide history, and subjective sense of adaptive functioning. Emotion dysregulation added significant incremental validity in relation to a broad range of distressful psychological symptoms and maladaptive behaviors.

The deleterious effects of early-life traumatic experiences appear to ripple through the lifespan as implicated in their relationships with current ratings of negative affect, posttraumatic stress, drug abuse, depression, and suicidality. Higher levels of childhood abuse and neglect were significantly related to higher levels of emotion dysregulation, adding support to theoretical models in which the developmental capacity to adaptively regulate emotions may be disturbed by early disruptive experiences.⁹

In addition, the data from this study support the conceptualization of emotion dysregulation as a distinct construct, related to but not reducible to negative affect.²⁷ The data suggest that negative affect may be as strongly or more strongly related to some forms of psychopathology than emotion dysregulation (eg, depression) but not to others, particularly

more impulsive, self-destructive, or externalizing disorders and behaviors such as substance abuse and suicide attempts. Also, given the high levels of trauma exposure and daily life stress (eg, poverty, neighborhood violence) to which the population of this study was exposed, it is possible that both suicide attempts and substance abuse themselves represent maladaptive emotion regulation strategies that are more likely to be employed in the face of stress by those with higher levels of dysregulated emotions.

Limitations

The primary limitations of this study arise from the use of cross-sectional and retrospective data collection. Due to the cross-sectional nature of the data, we are unable to determine the degree to which emotion dysregulation is a risk factor for development of adult psychopathology, a consequence of certain forms of psychopathology, or a central component of various forms of psychopathology for which it is not a diagnostic criterion. For example, the combination of intrusive experiences, avoidance/numbing, and hyperarousal associated with PTSD could lead to an overall state of emotion dysregulation. Much of the literature on emotion dysregulation suggests a childhood etiology, often involving traumatic adverse childhood experiences including childhood abuse and lack of a secure attachment with caregivers.⁵⁶⁻⁵⁹ Supporting this model, some longitudinal research indicates that early emotion regulation problems predict later risk for psychopathology.⁶⁰⁻⁶²

Consistent with the data in this study, a number of recent studies^{5,63,64} also suggest that emotion dysregulation may be a “higher order” factor that cuts across multiple psychiatric disorders. Conversely, studies⁶⁵⁻⁶⁸ focused on heterogeneity within psychiatric diagnoses suggest that subtypes of multiple psychiatric disorders (eg, eating disorders, personality disorders, anxiety disorders, and mood disorders) may be associated with varying degrees of emotion dysregulation.⁶⁵⁻⁶⁸ Alternately, the relationship between emotion dysregulation and some psychological disorders may be an interactive cascade in which emotion dysregulation increases vulnerability for the development of psychiatric disorders that in turn exacerbate emotion dysregulation. In PTSD, the presence of trauma-related cues may lead to emotion dysregulation that in turn may lead to higher levels of PTSD symptoms such as avoidance and irritability.

Longitudinal studies are needed to examine the developmental relationship of early life stressors, emotion dysregulation, and psychopathology. In addition, more research evaluating the biomarkers (eg, genetic data, imaging data) of negative affect and emotion dysregulation, as well as the relationship between the 2 variables, is needed to understand the way in which these 2 traits develop, function biologically, interact, and contribute to varying levels and types of psychopathology.

Additionally, we had a relatively homogeneous sample with respect to both race and income, making it important to consider the possibility that factors specific to this low-income, urban, primarily African American sample may

relate to the results of the study. For example, a number of studies⁶⁹⁻⁷² have found a relationship between experiences of racial discrimination and the risk for psychological disorders. Other recent studies⁷³⁻⁷⁵ focusing on emotion dysregulation and related constructs (eg, distress tolerance) point to the importance of taking into account race, gender, and socioeconomic status; one of these recent studies⁷³ found that low levels of distress tolerance conferred increased risk for alcohol use among whites, delinquent behavior among African Americans, and internalizing symptoms among women. These types of findings highlight the importance of social context in psychiatric research and in the development of treatments and preventive intervention. At the same time, this sample represents a population at very high risk for both depression and PTSD that is generally underrepresented in research on psychopathology. Thus, the data have valuable potential for informing broader public health policies and practices.

Finally, as research on emotion dysregulation has increased dramatically over the last decade, a number of clinician/interviewer-rated and self-report instruments assessing emotion dysregulation have been developed that span a range of underlying constructs.^{76,77} The emotion dysregulation measurement used in this study was grounded in research on personality disorders, particularly borderline personality disorder,⁷⁸ and the developmental etiology of personality.⁷⁹ Our measurement instrument was also based on prior work conducted by our research team on the assessment of affect regulation and dysregulation.^{15,27,42,43,78} As with other research in related areas including personality pathology and adult attachment styles, we expect that data gathered from this self-report instrument will diverge somewhat from interviewer-rated and clinician-rated instruments.⁸⁰⁻⁸² Ideally, future research would include both self-report instruments and clinician/interviewer-rated instruments, tapping multiple facets of emotion regulation. However, we hope that the Emotion Dysregulation Scale used in this study, which was designed on the basis of data from general psychiatric samples and is not intended for any single population, may serve as a useful instrument for the assessment of emotion dysregulation in at-risk or clinical populations.

Clinical Implications

Emotion dysregulation is believed to present in extremes of either overly restricted emotional expression and avoidance or heightened and excessive emotionality and excitement-seeking evident across a number of types of psychopathology.⁸³ Rumination, panic, self-criticism, social inhibition, interpersonal isolation, concentration difficulties, and attention problems may reflect internalized failures of emotion management, while the externalized behaviors such as aggression, alcohol and substance abuse, disordered eating, self-harm, and suicidality associated with emotion dysregulation may represent, in part, efforts to escape emotions experienced as overwhelming or intolerable.^{26,84-87}

Facets of emotion dysregulation may represent core clinical targets in treatments for a range of psychological

disorders, not just borderline personality disorder.⁸⁸ A greater understanding and more effective measurement of emotion dysregulation may facilitate the process of collaborative goal formation in psychotherapeutic treatments. Treatment goals in individuals presenting with emotion dysregulation might focus on the reduction of affective arousal; increase in affect tolerance; cognitive reappraisal of negative affect states, causes, and consequences; increase in social-perspective-taking during periods of affective intensity; and development of psychosocial coping skills to be used in times of increased emotional distress.

Disclosure of off-label usage: The authors have determined that, to the best of their knowledge, no investigational information about pharmaceutical agents that is outside US Food and Drug Administration-approved labeling has been presented in this article.

Author affiliations: Department of Psychiatry and Behavioral Sciences (Drs Bradley, Ressler, and Westen and Mss Phifer and Fani) and Department of Psychology (Drs DeFife, Guarnaccia, and Westen), Emory University School of Medicine, and Yerkes National Primate Research Center (Dr Ressler), Emory University, Atlanta, Georgia; Atlanta Veterans Affairs Medical Center, Atlanta, Georgia (Dr Bradley); and Howard Hughes Medical Institute, Chevy Chase, Maryland (Dr Ressler).

Financial disclosure: Drs Bradley, DeFife, Guarnaccia, Ressler, and Westen and Mss Phifer and Fani have no personal affiliations or financial relationships with any commercial interest to disclose relative to the article.

Funding/support: This work was primarily supported by the National Institute of Mental Health (grants MH071537, MH078100, and MH082256), Bethesda, Maryland; Emory and Grady Memorial Hospital General Clinical Research Center, Atlanta, Georgia; the National Institutes of Health National Centers for Research Resources (grant M01RR00039), Bethesda, Maryland; the American Foundation for Suicide Prevention, New York, New York; and the Burroughs Wellcome Fund, Research Triangle Park, North Carolina.

REFERENCES

- Watson D, Clark LA. Affects separable and inseparable: on the hierarchical arrangement of the negative affects. *J Pers Soc Psychol*. 1992;62(3):489–505.
- Krueger RF. The structure of common mental disorders. *Arch Gen Psychiatry*. 1999;56(10):921–926.
- Gross JJ, ed. *Handbook of Emotion Regulation*. New York, NY: Guilford Press; 2007.
- Linehan MM, Heard HL. Dialectical behavior therapy for borderline personality disorder. In: Clarkin JF, Marziali E, Munroe-Blum H, eds. *Borderline Personality Disorder: Clinical and Empirical Perspectives*. New York, NY: Guilford Press; 1992:248–267.
- Shedler J, Westen D. Dimensions of personality pathology: an alternative to the five-factor model. *Am J Psychiatry*. 2004;161(10):1743–1754.
- Westen D. Cultural, emotional, and unconscious aspects of self. In: Curtis RC, ed. *The Relational Self*. New York, NY: Guilford Press; 1991:181–210.
- Westen D. Affect regulation and psychopathology: applications to depression and borderline personality disorder. In: Flack W, Laird J, eds. *Affect and Psychopathology*. New York, NY: Oxford University Press; 1998.
- Cole PM, Michel MK, Teti LO. The development of emotion regulation and dysregulation: a clinical perspective. *Monogr Soc Res Child Dev*. 1994;59(2–3):73–100.
- Calkins SD, Hill A. Caregiver influences on emerging emotion regulation: biological and environmental transactions in early development. In: Gross JJ, ed. *Handbook of Emotion Regulation*. New York, NY: Guilford Press; 2007:229–248.
- De Pauw SS, Mervielde I. Temperament, personality and developmental psychopathology: a review based on the conceptual dimensions underlying childhood traits. *Child Psychiatry Hum Dev*. 2010;41(3):313–329.
- Krueger RF, Hicks BM, Patrick CJ, et al. Etiologic connections among substance dependence, antisocial behavior, and personality: modeling the externalizing spectrum. *J Abnorm Psychol*. 2002;111(3):411–424.
- Krueger RF, Tackett JL. Personality and psychopathology: working toward the bigger picture. *J Pers Disord*. 2003;17(2):109–128.
- Nigg JT. On inhibition/disinhibition in developmental psychopathology: views from cognitive and personality psychology and a working inhibition taxonomy. *Psychol Bull*. 2000;126(2):220–246.
- Anda RE, Felitti VJ, Bremner JD, et al. The enduring effects of abuse and related adverse experiences in childhood: a convergence of evidence from neurobiology and epidemiology. *Eur Arch Psychiatry Clin Neurosci*. 2006;256(3):174–186.
- Zittel Conklin C, Westen D. Borderline personality disorder in clinical practice. *Am J Psychiatry*. 2005;162(5):867–875.
- Bradley R, Jenei J, Westen D. Etiology of borderline personality disorder: disentangling the contributions of intercorrelated antecedents. *J Nerv Ment Dis*. 2005;193(1):24–31.
- Skodol AE, Siever LJ, Livesley WJ, et al. The borderline diagnosis II: biology, genetics, and clinical course. *Biol Psychiatry*. 2002;51(12):951–963.
- Trull TJ, Widiger TA, Lynam DR, et al. Borderline personality disorder from the perspective of general personality functioning. *J Abnorm Psychol*. 2003;112(2):193–202.
- Westen D, Heim AK, Morrison K, et al. Classifying and diagnosing psychopathology: a prototype matching approach. In: Beutler L, Malik M, eds. *Rethinking the DSM: A Psychological Perspective*. Washington, DC: American Psychological Association Press; 2002:221–250.
- Zanarini MC, Frankenburg FR, DeLuca CJ, et al. The pain of being borderline: dysphoric states specific to borderline personality disorder. *Harv Rev Psychiatry*. 1998;6(4):201–207.
- Zanarini MC, Frankenburg FR, Hennen J, et al. The longitudinal course of borderline psychopathology: 6-year prospective follow-up of the phenomenology of borderline personality disorder. *Am J Psychiatry*. 2003;160(2):274–283.
- Coccaro EF, Bergeman CS, McClearn GE. Heritability of irritable impulsiveness: a study of twins reared together and apart. *Psychiatry Res*. 1993;48(3):229–242.
- Sanislow CA, Grilo CM, McGlashan TH. Factor analysis of the DSM-III-R borderline personality disorder criteria in psychiatric inpatients. *Am J Psychiatry*. 2000;157(10):1629–1633.
- Sanislow CA, Grilo CM, Morey LC, et al. Confirmatory factor analysis of DSM-IV criteria for borderline personality disorder: findings from the Collaborative Longitudinal Personality Disorders Study. *Am J Psychiatry*. 2002;159(2):284–290.
- Zanarini MC, Frankenburg FR, Yong L, et al. Borderline psychopathology in the first-degree relatives of borderline and Axis II comparison probands. *J Pers Disord*. 2004;18(5):439–447.
- Yen S, Zlotnick C, Costello E. Affect regulation in women with borderline personality disorder traits. *J Nerv Ment Dis*. 2002;190(10):693–696.
- Westen D, Muderrisoglu S, Fowler C, et al. Affect regulation and affective experience: individual differences, group differences, and measurement using a Q-sort procedure. *J Consult Clin Psychol*. 1997;65(3):429–439.
- Linehan MM. *Cognitive-Behavioral Treatment of Borderline Personality Disorder*. New York, NY: Guilford Press; 1993.
- Kellogg SH, Young JE. Schema therapy for borderline personality disorder. *J Clin Psychol*. 2006;62(4):445–458.
- Young J, Klosko J, Weishaar M. *Schema Therapy: A Practitioner's Guide*. New York, NY: Guilford Press; 2003.
- Clarkin JF, Levy KN, Lenzenweger MF, et al. Evaluating three treatments for borderline personality disorder: a multiwave study. *Am J Psychiatry*. 2007;164(6):922–928.
- Miller JD, Pilkonis PA. Neuroticism and affective instability: the same or different? *Am J Psychiatry*. 2006;163(5):839–845.
- McDermott MJ, Tull MT, Gratz KL, et al. The role of anxiety sensitivity and difficulties in emotion regulation in posttraumatic stress disorder among crack/cocaine dependent patients in residential substance abuse treatment. *J Anxiety Disord*. 2009;23(5):591–599.
- Gillespie CF, Bradley B, Mercer K, et al. Trauma exposure and stress-related disorders in inner city primary care patients. *Gen Hosp Psychiatry*. 2009;31(6):505–514.
- Alim TN, Graves E, Mellman TA, et al. Trauma exposure, posttraumatic stress disorder and depression in an African-American primary care population. *J Natl Med Assoc*. 2006;98(10):1630–1636.
- Chapman DP, Whitfield CL, Felitti VJ, et al. Adverse childhood experiences and the risk of depressive disorders in adulthood. *J Affect Disord*. 2004;82(2):217–225.
- Binder EB, Bradley RG, Liu W, et al. Association of FKBP5 polymorphisms and childhood abuse with risk of posttraumatic stress disorder symptoms in adults. *JAMA*. 2008;299(11):1291–1305.
- Bradley RG, Binder EB, Epstein MP, et al. Influence of child abuse on adult depression: moderation by the corticotropin-releasing hormone receptor gene. *Arch Gen Psychiatry*. 2008;65(2):190–200.
- Bernstein D, Fink L. *Childhood Trauma Questionnaire Manual*. San Antonio, TX: Psychological Corporation; 1998.

40. Bernstein DP, Stein JA, Newcomb MD, et al. Development and validation of a brief screening version of the Childhood Trauma Questionnaire. *Child Abuse Negl.* 2003;27(2):169–190.
41. Watson D, Clark LA, Tellegen A. Development and validation of brief measures of positive and negative affect: the PANAS scales. *J Pers Soc Psychol.* 1988;54(6):1063–1070.
42. Westen D, Shedler J. Revising and assessing Axis II, part 1: developing a clinically and empirically valid assessment method. *Am J Psychiatry.* 1999;156(2):258–272.
43. Westen D, Shedler J. Revising and assessing Axis II, part 2: toward an empirically based and clinically useful classification of personality disorders. *Am J Psychiatry.* 1999;156(2):273–285.
44. Blake D, Weathers FW, Nagy LM, et al. A clinician rating scale for assessing current lifetime PTSD: the CAPS-1. *Behav Ther (N Y N Y).* 1990;13:187–188.
45. Weathers FW, Keane TM, Davidson JR. Clinician-Administered PTSD Scale: a review of the first ten years of research. *Depress Anxiety.* 2001;13(3):132–156.
46. Blake DD, Weathers FW, Nagy LM, et al. The development of a clinician administered PTSD scale. *J Trauma Stress.* 1995;8(1):75–90.
47. Saunders JB, Aasland OG, Babor TF, et al. Development of the Alcohol Use Disorders Identification Test (AUDIT): WHO collaborative project on early detection of persons with harmful alcohol consumption-II. *Addiction.* 1993;88(6):791–804.
48. Skinner HA. The Drug Abuse Screening Test. *Addict Behav.* 1982;7(4):363–371.
49. Cocco KM, Carey KB. Psychometric properties of the Drug Abuse Screening Test in psychiatric outpatients. *Psychol Assess.* 1998;10(4):408–414.
50. Maisto SA, Carey MP, Carey KB, et al. Use of the AUDIT and the DAST-10 to identify alcohol and drug use disorders among adults with a severe and persistent mental illness. *Psychol Assess.* 2000;12(2):186–192.
51. French MT, Roebuck MC, McGeary KA, et al. Using the Drug Abuse Screening Test (DAST-10) to analyze health services utilization and cost for substance users in a community-based setting. *Subst Use Misuse.* 2001;36(6–7):927–946.
52. Beck AT, Steer RA, Ball R, et al. Comparison of Beck Depression Inventories-IA and -II in psychiatric outpatients. *J Pers Assess.* 1996;67(3):588–597.
53. Beck AT, Steer RA, Brown GK. *Manual for the Beck Depression Inventory-II.* San Antonio, TX: Psychological Corporation; 1996.
54. Keller MB, Lavori PW, Friedman B, et al. The Longitudinal Interval Follow-up Evaluation: a comprehensive method for assessing outcome in prospective longitudinal studies. *Arch Gen Psychiatry.* 1987;44(6):540–548.
55. O'Brien R. A caution regarding rules of thumb for variance inflation factors. *Qual Quant.* 2007;41(5):673–690.
56. Riggs S. Childhood emotional abuse and the attachment system across the life cycle: what theory and research tell us. *J Aggress Maltreat Trauma.* 2010;19(1):5–51.
57. Kochanska G, Philibert RA, Barry RA. Interplay of genes and early mother-child relationship in the development of self-regulation from toddler to preschool age. *J Child Psychol Psychiatry.* 2009;50(11):1331–1338.
58. Alink LR, Cicchetti D, Kim J, et al. Mediating and moderating processes in the relation between maltreatment and psychopathology: mother-child relationship quality and emotion regulation. *J Abnorm Child Psychol.* 2009;37(6):831–843.
59. Cicchetti D, Ackerman B, Izard C. Emotions and emotion regulation in developmental psychopathology. *Dev Psychopathol.* 1995;7(1):1–10.
60. Kim J, Cicchetti D. Longitudinal pathways linking child maltreatment, emotion regulation, peer relations, and psychopathology. *J Child Psychol Psychiatry.* 2010;51(6):706–716.
61. Brotman MA, Schmajuk M, Rich BA, et al. Prevalence, clinical correlates, and longitudinal course of severe mood dysregulation in children. *Biol Psychiatry.* 2006;60(9):991–997.
62. Hatzenbuehler ML, McLaughlin KA, Nolen-Hoeksema S. Emotion regulation and internalizing symptoms in a longitudinal study of sexual minority and heterosexual adolescents. *J Child Psychol Psychiatry.* 2008;49(12):1270–1278.
63. Aldao A, Nolen-Hoeksema S, Schweizer S. Emotion-regulation strategies across psychopathology: a meta-analytic review. *Clin Psychol Rev.* 2010;30(2):217–237.
64. Taylor SF, Liberzon I. Neural correlates of emotion regulation in psychopathology. *Trends Cogn Sci.* 2007;11(10):413–418.
65. Jones M, Westen D. Diagnosis and subtypes of adolescent antisocial personality disorder. *J Pers Disord.* 2010;24(2):217–243.
66. Thompson-Brenner H, Eddy KT, Satir DA, et al. Personality subtypes in adolescents with eating disorders: validation of a classification approach. *J Child Psychol Psychiatry.* 2008;49(2):170–180.
67. Lanius RA, Vermetten E, Loewenstein RJ, et al. Emotion modulation in PTSD: Clinical and neurobiological evidence for a dissociative subtype. *Am J Psychiatry.* 2010;167(6):640–647.
68. Ortigo KM, Westen D, Bradley B. Personality subtypes of suicidal adults. *J Nerv Ment Dis.* 2009;197(9):687–694.
69. Williams DR, Neighbors HW, Jackson JS. Racial/ethnic discrimination and health: findings from community studies. *Am J Public Health.* 2008;98(suppl):S29–S37.
70. Himle JA, Baser RE, Taylor RJ, et al. Anxiety disorders among African Americans, blacks of Caribbean descent, and non-Hispanic whites in the United States. *J Anxiety Disord.* 2009;23(5):578–590.
71. Ellis BH, MacDonald HZ, Lincoln AK, et al. Mental health of Somali adolescent refugees: the role of trauma, stress, and perceived discrimination. *J Consult Clin Psychol.* 2008;76(2):184–193.
72. Weems CF, Watts SE, Marsee MA, et al. The psychosocial impact of Hurricane Katrina: contextual differences in psychological symptoms, social support, and discrimination. *Behav Res Ther.* 2007;45(10):2295–2306.
73. Daughters SB, Reynolds EK, MacPherson L, et al. Distress tolerance and early adolescent externalizing and internalizing symptoms: the moderating role of gender and ethnicity. *Behav Res Ther.* 2009;47(3):198–205.
74. Bornovalova MA, Gratz KL, Daughters SB, et al. A multimodal assessment of the relationship between emotion dysregulation and borderline personality disorder among inner-city substance users in residential treatment. *J Psychiatr Res.* 2008;42(9):717–726.
75. Gratz KL, Tull MT, Baruch DE, et al. Factors associated with co-occurring borderline personality disorder among inner-city substance users: the roles of childhood maltreatment, negative affect intensity/reactivity, and emotion dysregulation. *Compr Psychiatry.* 2008;49(6):603–615.
76. Gross JJ, John OP. Individual differences in two emotion regulation processes: implications for affect, relationships, and well-being. *J Pers Soc Psychol.* 2003;85(2):348–362.
77. Gratz K, Roemer L. Multidimensional assessment of emotion regulation and dysregulation: development, factor structure, and initial validation of the Difficulties in Emotion Regulation Scale. *J Psychopathol Behav Assess.* 2004;26(1):41–54.
78. Conklin CZ, Bradley R, Westen D. Affect regulation in borderline personality disorder. *J Nerv Ment Dis.* 2006;194(2):69–77.
79. Bradley R, Westen D. The psychodynamics of borderline personality disorder: a view from developmental psychopathology. *Dev Psychopathol.* 2005;17(4):927–957.
80. Dozier M, Lee S. Discrepancies between self- and other-report of psychiatric symptomatology: effects of dismissing attachment strategies. *Dev Psychopathol.* 1995;7(1):217–226.
81. Bradley R, Hilsenroth M, Guarnaccia C, et al. Relationship between clinician assessment and self-assessment of personality disorders using the SWAP-200 and PAL. *Psychol Assess.* 2007;19(2):225–229.
82. DeFife J, Drill R, Nakash O, et al. Agreement between clinician and patient ratings of adaptive functioning and developmental history [published online ahead of print July 15, 2010]. *Am J Psychiatry.* 2010;167(12):1472–1478.
83. Gross JJ, Thompson RA. Emotion regulation: conceptual foundations. In: Gross JJ, ed. *Handbook of Emotion Regulation.* New York, NY: Guilford Press; 2007:3–25.
84. Brown MZ, Comtois KA, Linehan MM. Reasons for suicide attempts and nonsuicidal self-injury in women with borderline personality disorder. *J Abnorm Psychol.* 2002;111(1):198–202.
85. Kullgren G. Factors associated with completed suicide in borderline personality disorder. *J Nerv Ment Dis.* 1988;176(1):40–44.
86. Montgomery SA, Montgomery D, Baldwin D, et al. Intermittent 3-day depressions and suicidal behaviour. *Neuropsychobiology.* 1989;22(3):128–134.
87. Markey MA, Vander Wal JS. The role of emotional intelligence and negative affect in bulimic symptomatology. *Compr Psychiatry.* 2007;48(5):458–464.
88. Barlow DH, Allen LB, Choate ML. Toward a unified treatment for emotional disorders. *Behav Ther.* 2004;35(2):205–230.

For the CME Posttest for this article, see pages 729–730.