

It is illegal to post this copyrighted PDF on any website.

Behavioral Symptoms Improve Prediction Models of Mortality in Patients With Dementia

To the Editor: The recent cohort study by Cheng et al¹ that focused on mortality predictors in Taiwanese patients with dementia yielded interesting results. The long follow-up period is a strength of the study. The authors chose a prediction model in which comorbidities and demographic variables were included in the analysis. In the Cox regression model, the use of antipsychotic drugs was not associated with mortality (hazard ratio = 1.09; 95% CI, 0.98–1.22).¹ This result helps clarify the controversy regarding an association of antipsychotic use and mortality in patients with dementia. I agree that there is not enough evidence to support this association, as was stated in a systematic review of mine published last year.² The lack of association of antidepressants with mortality is also reassuring and in agreement with a previous report.³

Although the prediction model presented is useful for clinicians, I would have welcomed a model including any scale measuring behavioral disturbances (depression, agitation, aggression, hallucinations), which are highly prevalent and predictive of mortality.^{2–6} It would also be interesting to know the proportion of patients with Alzheimer’s type dementia and that of other types of dementia to compare the results with those of the cohorts from Western countries. It seems that vascular dementia is more frequent in Asia than in these countries.

REFERENCES

1. Cheng CM, Chang WH, Chiu YC, et al. Risk score for predicting mortality in people with dementia: a nationwide population-based cohort study in Taiwan with 11 years of follow-up. *J Clin Psychiatry*. 2019;80(4):18m12629.
2. Modrego PJ, Lobo A. Determinants of progression and mortality in Alzheimer’s disease: a systematic review. *Neuropsychiatry (London)*. 2018;8(5):1465–1475.
3. Lopez OL, Wisniewski SR, Becker JT, et al. Psychiatric medication and abnormal behavior as predictors of progression in probable Alzheimer disease. *Arch Neurol*. 1999;56(10):1266–1272.
4. Neumann PJ, Araki SS, Arcelus A, et al. Measuring Alzheimer’s disease progression with transition probabilities: estimates from CERAD. *Neurology*. 2001;57(6):957–964.
5. Tun SM, Murman DL, Long HL, et al. Predictive validity of neuropsychiatric subgroups on nursing home placement and survival in patients with Alzheimer disease. *Am J Geriatr Psychiatry*. 2007;15(4):314–327.
6. Magni E, Binetti G, Bianchetti A, et al. Risk of mortality and institutionalization in demented patients with delusions. *J Geriatr Psychiatry Neurol*. 1996;9(3):123–126.

Pedro J. Modrego, MD^a
pmodrego@salud.aragon.es

^aDepartment of Neurology, Hospital Miguel Servet, Saragoza, Spain

Published online: March 24, 2020.

Potential conflicts of interest: None.

Funding/support: None.

J Clin Psychiatry 2020;81(3):19lr13217

To cite: Modrego PJ. Behavioral symptoms improve prediction models of mortality in patients with dementia. *J Clin Psychiatry*. 2020;81(3):19lr13217.

To share: <https://doi.org/10.4088/JCP.19lr13217>

© Copyright 2020 Physicians Postgraduate Press, Inc.

You are prohibited from making this PDF publicly available.

Dr Cheng and Colleagues Reply

To the Editor: We thank Dr Modrego for the response to our article.¹ We acknowledge that Drs Modrego and Lobo, in their systematic review,² stated that there is still not enough evidence to conclude that a link exists between antipsychotic use and mortality in dementia patients. Confounding factors may limit the results of observational studies investigating this question, but randomized controlled trials may fail to reflect real-world clinical situations because of their inclusion and exclusion criteria.³ A similar limitation also exists for investigations of the association between antidepressant use and mortality in dementia patients. To date, findings regarding the impact of antidepressants on mortality in dementia patients have been inconsistent.^{1,4-6}

On the other hand, the presence of behavioral and psychological symptoms of dementia (BPSD) may be a valuable factor in predicting mortality in dementia patients.² For example, agitation and aggression may cause head trauma or fractures due to falls and therefore increase the risk of death.⁷ Lopez et al⁷ found that the increased risk of death in patients with probable Alzheimer's disease was associated with the presence of psychiatric symptoms rather than exposure to antipsychotics. Our registry database, the National Health Insurance Research Database, did not provide information about BPSD or neuropsychiatric inventory results. Consequently, we could not evaluate the influence of these factors. However, the severity and type of BPSD may change over time.⁸ BPSD symptoms may elevate the risk of falls and fracture events, which were evaluated as the factor "femoral neck fracture" in our article.¹

The proportions of patients with Alzheimer's type dementia (ICD-9 code = 331.0) and vascular type dementia (ICD-9 code = 290.4) in the derivation cohort were 9.9% and 15.9%, respectively. We performed another analysis to add those factors of different dementia types into our Cox proportional hazards regression model, which was equal to the creating process described in our article.¹ The results showed that neither Alzheimer's type nor vascular type dementia was significant as a factor for the final variables selection (vascular type: $P = .79$, Alzheimer's type: $P = .974$) compared with other predicting variables. This result was generally in accord with previous findings. In a brain neuropathology study, the authors found few dementia cases that did not have a mixed component of both Alzheimer's-type pathologies and vascular lesions.⁹ According to the literature, although vascular dementia may be associated with a slightly higher mortality rate, the excess mortality in patients with certain stroke features may be driven greatly by cardiovascular disease outside the brain.^{10,11} Stroke or vascular dementia may demonstrate a phenomenon of worse generalized vascular disease that is not specific to the brain. More investigations will be required to clarify if type of dementia (especially vascular and Alzheimer's type dementia) impacts significantly as an independent predictor in the prediction model with adjustment for comorbidity variables (ie, competing mortality risks).

REFERENCES

1. Cheng CM, Chang WH, Chiu YC, et al. Risk score for predicting mortality in people with dementia: a nationwide, population-based cohort study in

- Taiwan with 11 years of follow-up. *J Clin Psychiatry*. 2019;80(4):18m12629.
2. Modrego PJ, Lobo A. Determinants of progression and mortality in Alzheimer's disease: a systematic review. *Neuropsychiatry (London)*. 2018;8(5):1465-1475.
3. Yeh TC, Tzeng NS, Li JC, et al. Mortality risk of atypical antipsychotics for behavioral and psychological symptoms of dementia: a meta-analysis, meta-regression, and trial sequential analysis of randomized controlled trials. *J Clin Psychopharmacol*. 2019;39(5):472-478.
4. Su JA, Chang CC, Wang HM, et al. Antidepressant treatment and mortality risk in patients with dementia and depression: a nationwide population cohort study in Taiwan. *Ther Adv Chronic Dis*. 2019;10:2040622319853719.
5. Mueller C, Huntley J, Stubbs B, et al. Associations of neuropsychiatric symptoms and antidepressant prescription with survival in Alzheimer's disease. *J Am Med Dir Assoc*. 2017;18(12):1076-1081.
6. Jennum P, Baandrup L, Ibsen R, et al. Increased all-cause mortality with use of psychotropic medication in dementia patients and controls: a population-based register study. *Eur Neuropsychopharmacol*. 2015;25(11):1906-1913.
7. Lopez OL, Becker JT, Chang YF, et al. The long-term effects of conventional and atypical antipsychotics in patients with probable Alzheimer's disease. *Am J Psychiatry*. 2013;170(9):1051-1058.
8. Tible OP, Riese F, Savaskan E, et al. Best practice in the management of behavioural and psychological symptoms of dementia. *Ther Adv Neurol Disorder*. 2017;10(8):297-309.
9. Fernando MS, Ince PG; MRC Cognitive Function and Ageing Neuropathology Study Group. Vascular pathologies and cognition in a population-based cohort of elderly people. *J Neurol Sci*. 2004;226(1-2):13-17.
10. Garcia-Ptacek S, Farahmand B, Kåreholt I, et al. Mortality risk after dementia diagnosis by dementia type and underlying factors: a cohort of 15,209 patients based on the Swedish Dementia Registry. *J Alzheimers Dis*. 2014;41(2):467-477.
11. Knopman DS, Rocca WA, Cha RH, et al. Survival study of vascular dementia in Rochester, Minnesota. *Arch Neurol*. 2003;60(1):85-90.

Chih-Ming Cheng, MD^{a,b,c,d}
Mu-Hong Chen, MD, PhD^{a,b,c}
Cheng-Hung Yang, MD^{a,b}
Shih-Jen Tsai, MD^{a,b}
Chia-Fen Tsai, MD, PhD^{a,b,c}
 cftsai@vghtpe.gov.tw

^aDepartment of Psychiatry, Taipei Veterans General Hospital, Taipei, Taiwan

^bDivision of Psychiatry, School of Medicine, National Yang-Ming University, Taipei, Taiwan

^cInstitute of Brain Science, National Yang-Ming University, Taipei, Taiwan

^dDepartment of Psychiatry, Taipei Veterans General Hospital, Yuanshan Branch, Yilan, Taiwan

Published online: March 24, 2020.

Potential conflicts of interest: None.

Funding/support: The authors' study discussed in this letter was supported by grants from Taipei Veterans General Hospital (V105D10-001-MY2-2), Ministry of Science and Technology (MOST 107-2634-F-075-002), and Taipei Veterans General Hospital-National Yang-Ming University Excellent Physician Scientists Cultivation Program (No.106-V-B-084).

Role of the sponsor: None of the supporters of the study had any role in the study design, data collection, analysis, interpretation of results, writing of the report, or the ultimate decision to submit the manuscript for publication.

J Clin Psychiatry 2020;81(3):191r13217a

To cite: Cheng CM, Chen MH, Yang CH, et al. Dr Cheng and colleagues reply.

J Clin Psychiatry. 2020;81(3):191r13217a.

To share: <https://doi.org/10.4088/JCP.191r13217a>

© Copyright 2020 Physicians Postgraduate Press, Inc.