



Correction

In the article “Clinical and Cognitive Correlates of Suicide Attempts in Bipolar Disorder: Is Suicide Predictable?” by Alison M. Gilbert, PhD, and colleagues in the August 2011 issue (*J Clin Psychiatry* 2011;72[8]:1027–1033), the odds ratios in Table 3 have been corrected to:

Table 3. Logistic Regression Analyses of Clinical and Demographic Predictors of Suicide Attempt

Predictor	Odds Ratio	95% CI		β	SE	P^a
		Lower	Upper			
Age	0.98	0.90	1.06	-0.02	0.04	.58
Sex	0.48	0.42	10.41	0.74	0.82	.36
Race	0.95	0.13	6.65	-0.06	1.00	.96
Mood type	3.11	0.79	12.20	1.14	0.70	.10
Psychosis	0.04	0.00	1.52	-3.21	1.85	.08
Trauma history	0.28	0.04	2.11	-1.28	1.03	.22
BIS-II total score	0.99	0.93	1.06	-0.01	0.03	.83
Aggression Questionnaire	0.99	0.95	1.03	-0.01	0.02	.64
RFL total score	0.98	0.96	1.00	-0.02	0.01	.10
HDRS-17	0.90	0.79	1.03	-0.10	0.07	.12
CARS-M	1.02	0.86	1.20	0.02	0.09	.85
WRAT-III	0.93	0.84	1.03	-0.07	0.05	.16

^aAll P values are nonsignificant.

Abbreviations: BIS-II = Barratt Impulsiveness Scale-version II, CARS-M = Clinician-Administered Rating Scale for Mania, HDRS-17 = 17-item Hamilton Depression Rating Scale, RFL = Reason for Living Inventory, SE = standard error, WRAT-III = Wide Range Achievement Test reading subscale.

The online article at PSYCHIATRIST.COM has been corrected.